

Low energy bounds on Poincaré violation in causal set theory

Nemanja Kaloper and David Mattingly

Department of Physics, University of California, Davis, CA 95616

In the causal set approach to quantum gravity, Poincaré symmetry is modified by swerving in spacetime, induced by the random lattice discretization of the space-time structure. The broken translational symmetry at short distances is argued to lead to a residual diffusion in momentum space, whereby a particle can acquire energy and momentum by drift along its mass shell and a system in equilibrium can spontaneously heat up. We consider bounds on the rate of momentum space diffusion coming from astrophysical molecular clouds, nuclear stability and cosmological neutrino background. We find that the strongest limits come from relic neutrinos, which we estimate to constrain the momentum space diffusion constant by $k < 10^{-61} \text{ GeV}^3$ for neutrinos with masses $m_\nu > 0.01 \text{ eV}$, improving the previously quoted bounds by roughly 17 orders of magnitude.

PACS numbers: 11.25.-w, 11.25.Mj, 98.80.Cq, 98.80.Qc

astro-ph/0607485

I. INTRODUCTION

The construction of a complete theory of quantum gravity has been a Holy Grail of theoretical physics over the past decades. The quest for quantum gravity faces new challenges that go beyond the problems encountered so far when quantizing gauge theories. Theoretically, we are not even sure how to properly define the observables of quantum gravity, let alone construct the full dynamics in a background-independent way. Experimentally, the feebleness of gravitational force, which may remain weak all the way up to the $4D$ Planck scale, $m_{\text{Pl}} \sim 10^{19} \text{ GeV}$, does not give much hope that direct experimentation could shed more light on the nature of quantum gravity in the foreseeable future ¹.

Under the looming shadow of conceptual difficulties and without experimental beacons, one may still pursue a ‘bottom-up’ approach: one may at least test theoretical proposals for compatibility with current observations, and constrain the parameters which control new physics signifying the various short-distance completions. For example, extra dimensions [1] or Planck scale modifications of Lorentz invariance [2] might both yield experimentally testable phenomena at low energies that can differentiate between approaches and lead to useful bounds on the underlying theoretical frameworks. There may also be other interesting phenomena, yet to be fully explored.

In this note we will focus on one such novel phenomenon in causal set theory [3, 4, 5], dubbed the ‘swerve’ effect [6]. In causal set theory, a particle will hop along a random lattice, spontaneously acquiring energy-momentum by drifting along its mass shell, at a rate controlled by a dimensional parameter that ought to arise from the underlying microscopic framework. Although this diffusion of particle population’s momentum distribution

is Lorentz invariant, the full Poincaré symmetry arises only statistically, because the translational symmetries are violated by the swerves. As a consequence, a gas of particles will heat up ‘spontaneously’ over time. Clearly the rate of any such diffusion processes in momentum space must be bounded to avoid conflicts with observed properties of isolated systems in equilibrium, which appear largely consistent with energy conservation.

Here we explore these bounds on the scale of energy-momentum from the swerves. We consider the possibility that the swerves may 1) heat up old, very cold molecular clouds in our galaxy, 2) facilitate nuclear α -decay, by providing a channel whereby α -particles in a nucleus gain more kinetic energy and escape more easily, and 3) render the background relic neutrino density hotter than is allowed by cosmological bounds on hot dark matter.

We find that molecular cloud heating and α -decay rates yield bounds which are numerically comparable to the hydrogen heat-up limits of [6]. However, the bounds from the background relic neutrino density, that behaves as hot dark matter, yield much stronger constraints which we estimate to restrict the momentum diffusion coefficient to $k \lesssim 10^{-61} \text{ GeV}^3$, exceeding those of [6] by about 17 orders of magnitude. By noting that different processes occur at different energy scales, we can also weakly constrain the dependence of k on energy. In what follows, we briefly review the basic tenets of causal set theory and the spacetime swerves in section II, outline the processes that lead to heating of nonrelativistic gases in section III, and discuss the bounds in section IV. We set $\hbar = c = k_B = 1$ throughout.

II. CAUSAL SETS FOR PEDESTRIANS

We start with a brief review of causal set theory, for a more complete discussion see [7]. A causal set C is a partially ordered set (a poset), consisting of ‘points’ $x, y, \dots$ and relations $x \prec y$ which encode causal ordering. For $x \prec y$ one says that x is to the past of y . Further, the ordering obeys two more rules, *i)* $x \prec y, y \prec z \Rightarrow x \prec z$ and *ii)* $x \not\prec x$. The former is transitivity, establishing

¹ An exception may be that the observed weakness of gravity is just a mirage, generated by the presence of as yet undiscovered large dimensions [1], in which case the full experimentally accessible quantum gravity might lurk near TeV.

links between points: if y is in the past of z , and x in the past of y , then x is also in the past of z . The latter forbids closed causal curves. In general one takes a causal set to be finite, in the sense that for any ordered pair $\{x, y\}$, such that $x \prec y$, there are a finite number of points z in between, such that $x \prec z$ and $z \prec y$.

To relate causal set ideas with low energy physics, we need to retrieve a geometric structure from the underlying causal set by some coarse graining. One can do so by reverse-engineering, using the process of ‘sprinkling’, or a random selection of points from a manifold M , such that the probability of picking any point from a region depends only on the 4-volume V of the region. Such sets of points and inherited causal relations can be thought of as causal sets C_M which approximate M as random lattices on it. Then the coarse-grained information about the topology of M , its metric, dimension, etc. can be deduced from operators defined strictly on C_M , and so C_M contains all of the necessary geometric information at this level of precision.

In contrast to conventional discrete lattices with a fixed lattice spacing, which explicitly break Lorentz symmetry, it has been argued that random lattices in causal set theory violate Lorentz symmetry much more mildly. Since the lattice is random and distributed in the manifold according to the 4-volume of spacetime regions, which is a Lorentz invariant, the symmetry is preserved to a much higher degree, and should be recovered in the limit when the coarse graining over the causal set becomes large.

On the other hand, it has been hypothesized by [6] that because a causal set is discrete, translation invariance of the Poincaré symmetry and hence energy-momentum conservation ought to be modified. An irregular lattice underlying the macroscopic spacetime structure won’t even preserve a discrete subgroup of translations, and hence the dual space lattice will also be jagged, allowing diffusion along a given mass shell instead of an evolution with a conserved pseudo-momentum, as in the case of a regular lattice. Indeed, imagine a causal set C_M approximating Minkowski space, and consider a classical particle initially at rest at a point x in C_M . Its world line is a set of measure zero, and thus for irregular C_M there won’t be any points directly to the future of x . By the arrow of time the particle will move forward in time, but having no lattice point in the future of x it must ‘swerve’ slightly to reach the next point in the causal set. This will endow it with momentum and kinetic energy, which will therefore not be conserved. The resulting behavior is Brownian drift in momentum space. For a large population of classical particles at low energies it should be described by the diffusion equation

$$\frac{\partial \rho}{\partial \tau} = k \nabla_P^2 \rho - m^{-1} p^\mu \partial_\mu \rho. \quad (1)$$

Here ∇_P^2 is the invariant Laplacian on the mass shell of the particle, describing only the drift along mass shell, m is the mass, τ is the proper time, $\partial_\mu = \frac{\partial}{\partial x^\mu}$ is coordinate space derivative and ρ is the momentum distribution of

the population. The constant k is a diffusion coefficient that sets the rate of diffusion. Although (1) is Poincaré invariant per se, given the statistical interpretation of the particle momentum, the spreading of ρ will yield a change in the average magnitude of particle momenta at low energies. Thus the standard Poincaré symmetry and momentum conservation may be violated statistically.

While the naive ‘swerve’ picture that underlies (1) requires modifications at high energies, (1) is singled out as the unique Lorentz invariant diffusion equation for a population ρ , at the lowest order in momentum derivatives. Indeed, it reflects the condition that a coarse grained description of a Lorentz-invariant momentum space random motion also respects Lorentz invariance. Therefore one can argue that (1) is a rather generic prediction from causal sets, instead of being tied to some particular underlying model of particle propagation [6].

When energy is bounded below, diffusion will inevitably yield an increase of average energy and of kinetic energy fluctuations in large samples of particles. At times short compared to some characteristic time that controls this classical spreading of the particle energy, the standard picture based on fully Poincaré invariant physics should remain valid. Thus although one doesn’t know how to formulate a full quantum-mechanical description of swerving, which may have to await the advent of a complete quantum gravity theory based on causal sets, at short enough time scales one should be able to separate classical and quantum behavior from each other by virtue of the Ehrenfest’s theorem and simply focus only on the swerve-induced corrections to the leading-order standard dynamics. The leading order contribution of swerves to low energy physics phenomena should appear from the widening of the momentum distribution, inducing a weak, slow, energy-momentum non-conservation. Over long times, the aggregate effect of diffusion may interfere with the standard description and yield observable signatures, and bounds, on the rate of momentum diffusion.

Such a picture should hold even for agglomerates of particles bound together, when describing the motion of the center of mass. Indeed, consider a system of two particles bound together by some potential interaction. In the momentum space, each particle can be viewed as a point on mass shell, and the interaction as a ‘strut’ linking them together. If the free particles would swerve, so should this bound state, as the underlying nature of spacetime is not affected by the interaction. However, the problem of finding the allowed lattice points to which the particles can jump under time evolution will be complicated because the particles need to jump close enough in order not to rip up the strut linking them. Sometimes, this may turn out to be impossible, and the swerves may break up the bond. When the momentum lattice is dense, with a mean energy spacing of lattice points much smaller than the scale of the bond, such destructive jumps will be very rare, while in the description of non-destructive jumps, whose particle energy differential due to swerving

is smaller than the binding energy, the resulting picture should be similar to what one expects for a single elementary state.

The momentum diffusion parameter k has dimensions of energy cubed, so one would naturally expect it to be some convolution of the energy scale of the problem and some short distance cutoff that emerges from causal set theory. To define this scale, let V_C be the 4-volume for which on average there is one point inside, and define the ‘fundamental’ energy scale $E_C = V_C^{-1/4}$. This scale is the natural analogue of the conventional field theory UV cutoff, characterizing the momentum space dual lattice, and may be high as the 4D Planck scale m_{Pl} . Thus we expect that $k = c\varepsilon^{3-\gamma}E_C^\gamma$ is a reasonable parameterization of the diffusion coefficient, where c is a dimensionless number and ε is the mass scale of the problem, such that $p^2 = \varepsilon^2$. Hence ε is invariant on the mass shell, behaving as the effective mass, in agreement with the microscopic Poincaré invariance of (1). Further, in an effective field theory (EFT) framework one might normally expect that $c \sim \mathcal{O}(1)$, unless it is protected by some softly broken symmetry. In causal set theory, the broken translational symmetries may be protecting the smallness of c , if there is a limit in which they are restored and swerving completely extinguished. In such a case c might turn out to be smaller than the naive arguments suggest. One may wonder if it is even possible to design an EFT framework for swerving, but as long as swerving is very small, there should exist an approximate description that may befit. As far as we can tell, proving this is presently an open question, which we will not pursue here. Now, if γ were large and negative, k would have rapidly vanished when $\varepsilon \ll E_C$ and so there would have been no diffusion to speak of. On the other hand, the variance of a population from Brownian motion is a function of the number of steps taken. If a particle travels a distance L then as E_C increases the number of steps increases, allowing for more diffusion. Thus one expects γ to be non-negative², possibly allowing interesting diffusion phenomena as time goes on. In fact, if there existed some kind of effective field theory description of swerves, one would have thought that the most likely value of γ were three, because this would correspond to describing the violation of translational symmetry by a marginal operator, which gives the strongest effects at low energies, that still respects decoupling.

With this in mind, we now turn to the quantitative constraints on k . We will derive various bounds on k from the temperature history of astrophysical molecular clouds, α -decay, and most importantly the cosmic neutrino background, which will enable us to constrain both the absolute value of k and, to a degree, the power γ .

² If the usual dimensional analysis applies to the determination of k , one would expect $\gamma = 3$. Keeping γ arbitrary we allow for possible anomalous quantum effects.

III. SWERVING GASES

We will be interested in non-relativistic particles, since for very relativistic ones the shift of the mean energy of the distribution becomes strongly suppressed by the boost factors and slows down [8]. This can be glimpsed at by multiplying (1) by a power of m , assuming continuity of ρ and taking the limit $m \rightarrow 0$. This leaves $p^\mu \partial_\mu \rho \rightarrow 0$, which for a homogeneous and isotropic distribution yields $\partial_t \rho \rightarrow 0$. On the other hand, for a non-relativistic population the proper time τ reduces to the coordinate time t , and the evolution equation (1) reduces to [6]

$$\frac{\partial \rho}{\partial t} = k \nabla_P^2 \rho, \quad (2)$$

where ∇_P^2 is now the standard spatial Laplacian operator on momentum space $\mathbb{R}^3$. This follows because the non-relativistic limit $m \gg p$ really corresponds to the restriction of the dynamics to a region about the mass shell minimum of size $p \ll m$, where the mass shell degenerates to a 3D Euclidean momentum space.

The general causal solution to (2) can be derived using standard Green’s function techniques [9]. The time dependent solution for $\rho(p, t)$ with initial conditions $\rho(p, 0) = f(p)$ is given by

$$\begin{aligned} \rho(p, t) &= \int_{-\infty}^{\infty} G(p, p', t, 0) f(p') d^3 p' \\ &= \int_{-\infty}^{\infty} (4\pi k t)^{-3/2} e^{-\frac{(p-p')^2}{4kt}} f(p') d^3 p', \end{aligned} \quad (3)$$

where $G(p, p', t, 0)$ is the Green’s function for the diffusion equation. One can immediately see the heating effect of swerves in a population by considering a collection of particles all initially at rest, $f(p') = \delta^3(p')$. In this case the integrals over $d^3 p'$ can immediately be done and one has simply

$$\rho(p) = (4\pi k t)^{-3/2} e^{-\frac{p^2}{4kt}}, \quad (4)$$

which is the thermal distribution for a non-relativistic gas at temperature $T = 2kt/m$, as mentioned in [6].

However in what follows we will consider the initial distribution of a population which is not at rest but is a “gas” at an initial temperature T_0 . This is consistent with the picture where the heating up from swerves is slow compared to the processes governed by conventional physics. Thus we need a solution that describes a system that starts with the standard Maxwell-Boltzman distribution $f(p') = (2\pi m T_0)^{-3/2} e^{-E'/T_0}$, and evolves by swerving into

$$\rho(p, t) = \int_{-\infty}^{\infty} (8\pi^2 k t m T_0)^{-3/2} e^{-\frac{(p-p')^2}{4kt}} e^{-\frac{p'^2}{2m T_0}} d^3 p'. \quad (5)$$

Factoring the constant part out and reducing the remaining integral to a Gaussian by completing the square in the exponent, one finds

$$\rho(p, t) = (2\pi m T)^{-3/2} e^{-p^2/(2m T)}, \quad (6)$$

where $T = T_0 + 2kt/m$. Hence a gas initially in thermal equilibrium will retain the same distribution under the influence of swerves. The temperature, however, increases linearly with time according to

$$\frac{dT}{dt} = \frac{2k}{m}. \quad (7)$$

Having thus established the primary effect of swerves in (6), (7), we can turn to exploring the bounds on k .

IV. BOUNDS

A. Nonrelativistic gases

From the nature of the swerve effect, one may expect that the strongest bounds will arise from the consideration of old and cold systems of light particles. Indeed, the bounds arise from limiting the gain of mean kinetic energy $\Delta\langle E \rangle \sim \Delta T$ due to swerves, which from (7) constrains k according to $k \lesssim m\Delta E/t$. Heavier particles will gain less energy due to their inertia, and clearly if the systems haven't existed for very long or if the energy thresholds are high, the bounds will be weak. In [6], the authors pursued this reasoning and limiting the rate of increase of temperature of a very cold hydrogen gas in normal laboratory conditions due to swerving by a millionth of a degree per second, they found the bound $k \lesssim 10^{-44} \text{ GeV}^3$.

One might hope to get better bounds from very old and cold astrophysical molecular clouds, which may be sensitive to heating by swerves. For example, in our galaxy there is a large molecular cloud called the Edge Cloud 2 [10] (EC2), at roughly 28 kpc from the galaxy center. It is known to contain several species of non-relativistic light molecules, including e.g. NH_3 , CO , and while it may even contain lighter molecules such as H_2 , to be conservative we will use here only those molecules which have been explicitly confirmed. Naively, this could yield a stronger bound³ than lab heating limits of [6]: limiting the temperature gain due to swerves by the average kinetic temperature of the NH_3 molecules in EC2, $T \simeq 20$ Kelvin, taking its estimated age as comparable to the age of the Milky Way, and using the molecular mass of approximately 17 GeV for NH_3 and finally and *crucially* assuming no dissipation, we would find a bound on k from (7) of $k \lesssim 10^{-51} \text{ GeV}^3$ that would appear to improve the hypothesized bound in [6] by about seven orders of magnitude.

However the assumption that dissipation is negligible is incorrect. The heat losses from electromagnetic emission reduce the above bound dramatically. The point is

that since the usual molecular clouds are composed of bound states of charges, they will behave as a system of elementary dipoles and emit electromagnetic waves when heated. Since the rate of energy gain from swerves is exceedingly small, and the clouds are not ultra-cold, the energy losses from electromagnetic radiation are faster and can neutralize the energy gain from swerves quickly, by a slight shift of the equilibrium temperature. Due to the uncertainties in determining this temperature, the quantitative bounds on k are weakened. To see this, let us consider a gas in equilibrium with its environment at a temperature T , determined to a small uncertainty $\Delta T \ll T$. At lowest order the uncertainty in the rate of energy loss due to radiation for a blackbody of surface area A , ignoring e.g. the temperature of the environment, such as CMB, with $c = \hbar = k_B = 1$ is

$$\Delta \frac{dE}{dt} \simeq \frac{4\pi^2}{60} AT^3 \Delta T. \quad (8)$$

Approximating $\frac{4\pi^2}{60} \sim 1$, and balancing this against the energy gain from swerves (7), $dE/dt = 3Nk/m$, where N is the number of particles, yields $AT^3 \Delta T = \frac{3Nk}{m}$. For a roughly spherical system, the lower limit on k that can be achieved by this is therefore

$$k \lesssim \frac{mT^3 \Delta T}{nR}, \quad (9)$$

where n is the number density and R is the linear dimension of the system. Using $n \sim 10^3$ particles per cubic centimeter [10], and a size on the order of a $R \sim \text{kpc}$ for EC2 the best limit we can achieve on k due to radiative emission is then

$$k \lesssim 10^{-41} \frac{\Delta T}{1 \text{ Kelvin}} \text{ GeV}^3. \quad (10)$$

The uncertainties in temperature for EC2 are within an order of magnitude of 1 Kelvin and so the limits on k really are $k \leq 10^{-41} \text{ GeV}^3$. In reality, this bound will be better when we properly include the grey body factors due to the material nature of emitting gases, that would obstruct the emission of electromagnetic waves, improving the bound by, at most, a couple of orders of magnitude and making it marginally comparable with the bounds claimed in [6]. We won't pursue those details here explicitly, because in what follows we will find a considerably stronger bound from the cosmological relic neutrino background.

This discussion however reveals that the problem of radiative losses will plague most attempts to constrain swerves. For example, one might try to use the atmospheres of Sun and Earth, which are composed of light gases and have been around for about five billion years⁴, and whose mean temperatures should not have varied by

³ Given that molecular clouds are not in exact thermal equilibrium and can even have strong localized deviations, these results must be considered approximate.

⁴ According, at least, to scientific sources.

more than, say, $\mathcal{O}(1)$. However as Sun and Earth equilibrate by radiative energy transfer, the bounds on k that one could get from constraining atmospheric temperature variation do not improve tremendously the numbers we quoted so far. Some of the problems with radiative losses could be mitigated by considering ultra-cold materials, for which the emission of electromagnetic waves, proportional to T^4 , will be dramatically reduced. For example, atom chip Bose-Einstein condensate (BEC) experiments with 87-Rubidium have been able to create a BEC with temperatures below a μ Kelvin that contain 2000 atoms in a volume $V \approx 10^{-3} \text{ mm}^3$ [11]. Using these parameters we find that we can neglect radiative losses if the diffusion parameter k is larger than $k \sim 10^{-54} \text{ GeV}^3$. On the other hand, the actual measured heating rate of these systems is $0.5 \mu\text{Kelvin}/\text{sec}$, giving a bound on k

$$k \lesssim 10^{-42} \text{ GeV}^3, \quad (11)$$

again within two orders of magnitude to the bound claimed in [6]. It may be of interest to carefully consider other such laboratory sources of bounds, however that is beyond the scope of the present work.

B. Atomic nuclei

Another example of very long-lived old systems of particles are atomic nuclei, and those which are particularly useful for our purposes here are the ones that undergo α -decay. As Gamow showed in 1928 [12], they may be viewed as systems comprised of a large number of α -particles which are bouncing around in the nuclear potential well. Their average energy controls the probability of tunneling out of the well, which causes the nucleus to decay. The whole process is well described by the stationary limit, where a particle of some typical kinetic energy $\sim \text{MeV}$ bounces many times against the barrier walls, occasionally managing to slip through it. If however an α particle managed to gain kinetic energy, it would move up inside the barrier, encountering thinner walls which would be easier to penetrate. Ultimately, a particle could even climb over the top of the wall and slip out unhindered.

In the standard Poincaré-invariant formulation of quantum mechanics, energy conservation implies that the energy levels of a bound state are discrete, as opposed to continuous as in a gas of free particles. One might therefore think that the α -particles in a nucleus should have discrete energy levels, and that the gap between them would prevent the swerving-induced energy gain of any individual particle, instead leading to only an overall gain of the center of mass energy of the whole nucleus. However, this intuition is misleading, because in causal set theory energy is not conserved absolutely. As a result, one would not get a system with fixed discrete levels, but with time-dependent energies which would ‘wobble’ around some statistical mean value, broadening the levels in a way similar to what happens in many body systems

at finite temperature. As a result, one would expect that a quantum state would really be represented by energy bands rather than sharp, discrete energy levels. Indeed, consider for example an α -particle in the ground eigenstate E_0 of a static unperturbed Hamiltonian H , where by unperturbed we mean without swerves. If we now include a small amount of swerving then under time evolution the particle cannot remain in E_0 since time translation is broken by the swerves. Since the swerves are random the evolution operator must have non-zero transition probabilities between states as there is certainly some probability for such a transition to happen all at once. Therefore the particles must evolve under swerves to a superposition of states with support on various eigenstates E_m of H , where $E_m > E_0$. The *mean value* of the α -particles would continuously gain energy, and we can estimate the rate of this energy gain by its classical value, appealing to Ehrenfest’s theorem. This should be a good approximation for deriving bounds on tunneling rates, by essentially saying that swerving enables tunneling when the broadening of individual energy levels is comparable to the gap between them in the Poincaré-invariant limit. This justifies the classical approximation in estimating energy gains in what follows.

Now, a typical stable heavy nucleus has a lifetime comparable to the age of the universe, 10^{10} years, and there are even some isotopes which are extremely long lived, such as Bi 209 whose lifetime is about 10^{19} years. On the other hand, the heavy nuclei which are present in nature are believed to have been created in first stars, and hence they are also very old, with their current ages comparable to their lifetimes. Because their measured lifetimes are in very good agreement with the α -decay theory based on the standard Poincaré invariant dynamics, we conclude that any possible enhancements of the decay due to swerving must be within the error of the measured lifetimes. This is why the longevity of Bi 209 does not help very much: since it is made artificially, it hasn’t been around for long enough to have swerves destabilize it.

From the functional dependence of the decay rate on the kinetic energy [13],

$$\Gamma \simeq \frac{AER}{(ER)^2 + \mathcal{B}}, \quad (12)$$

where $\mathcal{R}$ is the effective size of the nucleus and $\mathcal{A}, \mathcal{B}$ constants that depend on the barrier properties, the standard description will remain unaffected by swerves if the total energy gain of an α -particle inside the barrier remains smaller than a typical kinetic energy of an escaping α -particle. Roughly, this implies that we ought to constrain $\Delta E \lesssim \text{MeV}$ over the age of a nucleus, comparable to the age of the universe. That yields $k \lesssim m_\alpha \text{MeV} H_0$, and so we find

$$k_\alpha \lesssim 10^{-44} \text{ GeV}^3, \quad (13)$$

which, quite coincidentally, is again comparable with the hydrogen heating bound of [6]. Basically, for k smaller

than k_α , the diffusion in momentum space is too slow to help push the α -particle out of the nucleus. One might have hoped to gain a stronger bound from more stable isotopes. However, as we have noted above, being made in a lab, these are much younger than the age of the universe, and the swerves could not have had enough opportunity to build up a large ΔE yet.

C. Cosmic neutrinos

We have seen above that the most serious practical obstruction to constraining the swerve effect arises from radiative energy losses, that can overshadow the energy gain from swerves already at very low temperatures. This problem can be overcome if the thermal gas is composed of particles whose electromagnetic interactions are extremely suppressed. A perfect candidate for such a material are neutrinos. They are electrically neutral, and while they can have anomalous electromagnetic dipoles, generated by the quantum loops involving a lepton and a W , the relevant form-factors are very small. The conservative bounds on the magnetic dipole moment as quoted by the Particle Data Group [14]⁵ restrict the largest magnetic dipole moment, for ν_τ , by $\mu < 3.9 \times 10^{-7} \mu_B$, where μ_B is the Bohr magneton, and the electric dipole moment, again of ν_τ , by $d < 5.2 \times 10^{-17} \text{ ecm}$. For comparison, these are at least seven orders of magnitude smaller than the corresponding dipole moments for a hydrogen atom. Because dipole power losses are proportional to the square of the dipole moment, the emissivity of the neutrino gas will be at least fourteen orders of magnitude more suppressed than the emissivity of the hydrogen gas.

Hence we can treat cold neutrinos as an essentially non-radiating gas, which can therefore be a sensitive probe of any Poincaré-violating effects that can lead to spontaneous heat up. In what follows, we will consider implications of these observations for the cosmological neutrino background. We will show that swerving can generate enough heat in the cosmic neutrino background such that it would behave as hot dark matter and conflict with the observational evidence that dark matter should predominantly be cold. Requiring this does not happen will yield the strongest bound on the momentum space diffusion constant, constraining it to roughly $k \lesssim 10^{-61} \text{ GeV}^3$.

To start, we note that while the cosmological neutrino background has not been observed directly, given the successes of the Standard Model of particle physics and the Big Bang nucleosynthesis, there is little doubt that it exists (see, for example, a recent review articles [18, 19]). Furthermore, by the observed neutrino mass differences we believe that two neutrino species, at least, are heav-

ier than about 0.01 eV, with at least one species being heavier than 0.03 eV [20], while the third may be lighter, with a temperature slightly below the CMB temperature, $T_\nu \sim 10^{-4} \text{ eV}$. There may of course be even more light neutrinos, provided that they are Standard Model singlets. If they had masses in the 10^{-2} eV , they would strengthen our bounds even more.

These neutrinos were all in thermal equilibrium with the baryons and photons in the early universe, at temperatures well above neutrino masses. Because at those times the neutrinos were relativistic, the swerve effect was negligible for them, as we have pointed out above. As the universe expanded, the neutrino gas cooled and heavier neutrinos became nonrelativistic, so that eventually the swerving could reheat them again. Radiative energy losses could not compensate for this energy gain unless the diffusion parameter k were extremely small. To estimate this critical value of momentum diffusion coefficient k_{cr} induced by swerving, below which radiative losses become important, we can use a version of Eq. (9), with the emissivity factor $\xi < 10^{-14}$ included. Assuming that the heavier neutrino species have masses $m_\nu > 0.01 \text{ eV}$ [19], are distributed homogeneously inside the volume of the present horizon size $R \sim 1/H_0 \sim (10^{-33} \text{ eV})^{-1}$, with number density $n \sim T_\nu^3 \sim 10^{-12} \text{ eV}^3$ and uncertainty $\Delta T \sim T_\nu$, to be conservative, we find that the critical value of k_{cr} , below which radiative energy losses become important, is

$$k_{cr} \simeq \xi m_\nu T_\nu H_0 \simeq 10^{-80} \text{ GeV}^3. \quad (14)$$

If we were to consider these same neutrinos at the time when they froze out, at $T_\nu \sim m_\nu$, when $H_\nu \sim (T_\nu/T_0)^{3/2} H_0 \sim 1000 H_0$, we could raise k_{cr} by at most 5 orders of magnitude, which however remains negligibly small for our purposes here, as will be clear from the bound we are about to find. Thus, for all values of $k > k_{cr}$ we can completely ignore the electromagnetic emissions, and treat the background relic neutrinos as essentially an isolated system.

To understand what happens with the energy density of neutrinos with the swerving contributions included, in principle we would have to define, and solve, a cosmological version of the diffusion equation (1), that would also include the effects of cosmic expansion on the neutrino distribution and its mean energy. However, even without tackling this complicated problem, we can understand the qualitative features of the solution rather simply. Consider a thermal distribution of particles with number density $n \sim 1/a^3$ and mean energy per particle $\mathcal{E} \simeq m_\nu + T = m_\nu + T_0/a$, where we normalize $a_0 = 1$. Here, m is the rest mass of the particle species, whereas T is the mean kinetic energy per particle, or temperature, of the distribution. In the early universe, $T \gg m_\nu$, but as the universe expands T redshifts as $1/a$ and eventually becomes negligible compared to m_ν . Through this epoch, the energy density $\rho_\nu \sim \mathcal{E}n$ first scales as $\rho_\nu \sim T/a^3 \sim T_0/a^4$, eventually changing to $\rho_\nu \sim m_\nu/a^3$.

Enter the swerves. For a typical massive neutrino from

⁵ There may be even stronger bounds from astrophysical and cosmological considerations, some dating back to [15], and more in recent works [16, 17].

the distribution, the swerves begin to restore its kinetic energy at the rate $\partial T/\partial t \sim k/m_\nu$, as per Eq. (7). The swerving slows after the kinetic energy becomes relativistic again, reaching $T_* \gtrsim \text{few} \times m$, and slowly growing on. While the neutrinos are nonrelativistic, the mean energy gain per unit time is approximately constant, so we can estimate the time scale from $\partial T/\partial t \sim T_*/t_*$ and previous formulas, to be $t_* \sim m_\nu^2/k$. Thus after t_* , while the neutrino number density continues to dilute as $n \sim 1/a^3$, their characteristic energy per particle becomes relativistic again, $\mathcal{E} \gtrsim \text{few} \times m_\nu$. For a large enough k this energy continues to swerve up more slowly thereafter because the time scale of dilution by cosmic expansion is $\sim H^{-1}$, getting longer as the universe cools, so the swerving dominates, albeit it slows after particles regain relativistic speeds. The critical value of k for which enough energy is dumped back into the neutrinos to compensate for cosmic cooling can be easily approximated by equating the cooling and heating time scales ($H^{-1} \sim t_*$ in this case). As long as k is above this value then on average we will end up with a distribution of relativistic particles whose energy density dilutes only as $\rho_\nu \gtrsim \text{few} \times m_\nu/a^3$, instead of the usual $\rho \sim 1/a^4$ law.

Because massive neutrinos started out in equilibrium with other relativistic species in the universe, such as CMB, after t_* we can approximate their residual energy density per species as $\rho \gtrsim \text{few} \times m_\nu T^3$, and determine its fraction of the total relativistic matter content of the universe. Relative to the energy density of photons, $\rho_\gamma \sim T^4$, their fraction is

$$\Delta\omega_\nu = \frac{\rho_\nu}{\rho_\gamma} \gtrsim 10 \times \frac{m_\nu}{T}, \quad (15)$$

where we are using that there are at least 4 neutrino flavors more massive than $m_\nu \sim 0.01 \text{ eV}$, and that at least two are heavier than $3m_\nu \sim 0.03 \text{ eV}$, consistent with neutrino models as reviewed in [19, 20]. As the universe expands and T decreases, $\Delta\omega_\nu$ becomes larger, implying that the massive neutrino species contribute progressively more to the total relativistic energy density of the universe due to the injection of energy from the swerves. Thus as time goes on the relativistic fraction of the total energy density of the universe *increases*. This is similar to what would happen if there was some kind of late decay of nonrelativistic particles into relativistic matter, suddenly repopulating the universe with relativistic, hot dark matter.

The total amount of hot dark matter is bounded, implying that the maximal value of $\Delta\Omega_\nu = \Delta\omega_\nu \Omega_\gamma$ is bounded too. At very late times, this gives the inequality $\Delta\Omega_\nu < 0.2 \Omega_{DM}$ [21, 22]. These bounds on $\Delta\Omega_\nu$ translate into constraints on the rate of momentum diffusion induced by swerves. Indeed, suppose that the swerve-induced momentum diffusion constant k is large enough such that the time scale $t_* \sim m_\nu^2/k$ after which nonrelativistic neutrinos swerve back up to relativistic velocities is comparable to the age of the universe where the neutrino would have frozen out without swerves,

$$t_\nu \sim H_\nu^{-1} \sim (T_0/m_\nu)^{3/2} H_0^{-1}.$$

$$t_* \simeq t_\nu. \quad (16)$$

If this were the case, neutrinos with mass m_ν would never really cease to be relativistic, as we discussed above, since the redshift from cosmic expansion would be compensated by the swerving practically as soon as the temperature slides below m_ν . From this point onward, the neutrino species with masses m_ν would contribute relativistic energy density that would scale as $1/a^3$. Because $t_\nu \ll t_0$, today the effective neutrino fraction of the relativistic energy density would obey

$$\Delta\omega_\nu > 10 \times \frac{m_\nu}{T_0}, \quad (17)$$

or, plugging in the value $T_0 \sim 10^{-4} \text{ eV}$,

$$\Delta\omega_\nu > 10^5 \times \frac{m_\nu}{\text{eV}}, \quad (18)$$

which shows that for the neutrino masses $m_\nu > 0.01 \text{ eV}$, we would end up with $\Delta\omega_\nu > 10^3$. For larger k , $\Delta\omega_\nu$ would be even larger, due to continual energy buildup from slower swerving of relativistic particles. This would translate to an estimate of neutrino energy density in the present universe given by

$$\Delta\Omega_\nu = \Delta\omega_\nu \Omega_\gamma > 10^3 \Omega_\gamma \sim \Omega_{DM}. \quad (19)$$

which, being comparable with the cold dark matter density, would violate the existing bounds [21, 22, 23] by almost an order of magnitude, in the very least. In this case most of dark matter would be hot instead of cold. This implies that in fact the heat up of the neutrino gas by swerving should be reduced, until the time scale t_* is slower than t_ν . Requiring $t_* > t_\nu$ then yields

$$k < H_\nu m_\nu^2 \simeq \left(\frac{m_\nu}{T_0}\right)^{3/2} m_\nu^2 H_0. \quad (20)$$

Substituting $m_\nu \sim 0.01 \text{ eV}$, we finally obtain the estimate

$$k < 10^{-61} \text{ GeV}^3. \quad (21)$$

This improves the bounds from hydrogen heating up of [6] by about 17 orders of magnitude, and represents the principal result of this work. We stress that this bound is somewhat imprecise because we do not know the exact value of the neutrino mass and the details of the balancing act between cosmic expansion and swerving which sets the mean energy of the massive neutrino relics. However, one would not expect parametrically large changes of (21) from such corrections, as long as the neutrino masses are not extremely small. We also stress that the swerves controlled by a k obeying (21) would only impart minute effects on other particles. For example, a free electron could only gain kinetic energy $\Delta E < 0.1 \mu\text{eV}$ throughout the age of the universe, clearly a miniscule amount that would be easily masked away by thermal effects.

D. Scale dependence of momentum space diffusion

Now that we have constraints on k it is interesting to rewrite them in terms of our parameterization $k = c\varepsilon^{3-\gamma}E_C^\gamma$. The molecular masses that control the bounds for clouds, hydrogen and nuclear stability are $m \sim \text{few} \times m_{\text{proton}}$ where $m_{\text{proton}} \sim 1 \text{ GeV}$ is the proton mass, while the relevant mass scale for the cosmic neutrino bound is 0.01 eV . If we make the standard assumption that E_C , which represents the scale of the causal set, is the Planck energy then we can rewrite our parameterization as

$$\log c + \gamma(19 - \log \varepsilon) = \log \frac{k}{\varepsilon^3}, \quad (22)$$

where all energies are measured in GeV. Our limits on k , $k < 10^{-61} \text{ GeV}^3$ at $\varepsilon = 0.01 \text{ eV}$ and $k < 10^{-44} \text{ GeV}^3$ at $\varepsilon = 1 \text{ GeV}$ then exclude a two-dimensional region in $\gamma, \log c$ parameter space whose boundary are the lines

$$\log c + 19\gamma = -44 \text{ for } \gamma \leq \frac{16}{11}, \quad (23)$$

$$\log c + 30\gamma = -28 \text{ for } \gamma > \frac{16}{11}. \quad (24)$$

If we fix $\gamma = 3$, as suggested by EFT lore, then the corresponding constraint on c is $c < 10^{-118}$ which is an extremely small value begging for a first-principles explanation. Presenting a detailed and *precise* explanation of how c could become so small is a challenge before causal set theory. A possible way to argue it may happen it is to demonstrate that there is a limit where swerving is completely extinguished and so energy-momentum conservation and therefore translational symmetries are restored. In such an instance, one may believe that the coefficient c could be much smaller than unity even within the EFT description of swerving, if one existed, since the translational symmetries would then play a role of a softly broken symmetry [24], protecting c . Alternatively, perhaps causal set theory could never fully fit in the landscape of EFTs, in which case one must determine whatever it is that emerges in the low energy limit and replaces EFT.

Thus one still faces the problem of actually calculating k (or c) to determine precisely the link between its absurdly small value and the broken translational symmetries. This is reminiscent of the hierarchy problem in particle physics. From the causal set perspective, one could try to link this to the observation that if the sprinkling that models a random lattice is denser, characterized by a larger value of $E_C = V_C^{-1/4}$, then the probability of finding a lattice point near a mass shell anywhere

in spacetime increases, which might reduce swerving and lower the value of k . In the EFT language, this would suggest that k is *inversely* proportional to the ‘cutoff’ E_C , which would imply $\gamma < 0$ and effectively render k very small, and swerving irrelevant, at low energies. However such a detailed exploration of theoretical foundations of causal sets and the swerve effect is beyond the scope of the present work.

V. SUMMARY

In this article we have considered low energy bounds on momentum diffusion arising from swerves in causal set approach to quantum gravity. We find that the bounds coming from old and cold molecular clouds in our galaxy and nuclear stability are comparable to the bounds on the diffusion coefficient k coming from hydrogen heating up of [6]. However, we find that the bounds from the relic massive neutrino background coming from hot dark matter constraints are much stronger, leading to our estimate $k < 10^{-61} \text{ GeV}^3$ for neutrinos with masses $m_\nu > 0.01 \text{ eV}$, which would improve the bounds of [6] by about 17 orders of magnitude. We also note that if $\gamma = 3$, which is strongly favored by the standard effective field theory dimensional arguments, the swerve-induced effects must be tremendously suppressed for some as yet unknown reason. In this case the greatest effects of swerves would be on the light cosmological neutrinos, whose relic energy density would show surprising increase due to the energy injection from swerves. Oddly, if our bound is almost saturated, such effects could lend to some strange behavior in the equation of state of the universe. While such possibilities are amusing, exploring them in detail must await further developments in causal set theory.

Acknowledgments

We would like to thank M. Bucher, H.F. Dowker, L. Knox, K.A. Olive, R. Sorkin and especially M. Kaplinghat for useful discussions. We would also like to thank M. Kaplinghat for comments on the manuscript. NK would like to thank Galileo Galilei Institute, Florence, Italy, and to LPT, Universite de Paris-Sud, Orsay, France, for kind hospitality during the course of this work. NK and DM were supported in part by the DOE Grant DE-FG03-91ER40674. NK was also supported in part by the NSF Grant PHY-0332258 and by a Research Innovation Award from the Research Corporation.

[1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **429**, 263 (1998); Phys. Rev. D **59**, 086004 (1999); L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999); Phys. Rev. Lett. **83**, 4690 (1999).

[2] D. Mattingly, Living Rev. Rel. **8**, 5 (2005).

[3] L. Bombelli, J. H. Lee, D. Meyer and R. Sorkin, Phys. Rev. Lett. **59**, 521 (1987).

[4] R. D. Sorkin, Int. J. Theor. Phys. **36**, 2759 (1997).

- [5] M. Ahmed, S. Dodelson, P. B. Greene and R. Sorkin, Phys. Rev. D **69**, 103523 (2004).
- [6] F. Dowker, J. Henson and R. D. Sorkin, Mod. Phys. Lett. A **19**, 1829 (2004).
- [7] R. D. Sorkin, arXiv:gr-qc/0309009.
- [8] Swerving may still lead to the broadening of spectral lines of particles that hop about the lightcone, as argued in [6].
- [9] see, for example, the lecture notes by N. Andersson, <http://www.soton.ac.uk/~jhr/MA361/>.
- [10] D. A. Lubowich et al. in *Carnegie Observatories Astrophysics Series, Vol. 4: Origin and Evolution of the Elements*, 2003 ed. A. McWilliam and M. Rauch, <http://jupiter.phy.umist.ac.uk/~tjm/carnegie.pdf>
- [11] S. Du et al., Phys. Rev. A **70**, 053606 (2004).
- [12] G. Gamow, Zeitschrift für Physik, **51**, 204 (1928).
- [13] J.M. Blatt and V.F. Weisskopf, *Theoretical Nuclear Physics*, John Wiley & Sons, New York 1952.
- [14] S. Eidelman et al. [Particle Data Group], Phys. Lett. B **592**, 1 (2004).
- [15] J. Bernstein, M. Ruderman and G. Feinberg, Phys. Rev. **132**, 1227 (1963).
- [16] M. Fukugita and S. Yazaki, Phys. Rev. D **36**, 3817 (1987).
- [17] G. G. Raffelt, Phys. Rev. Lett. **64**, 2856 (1990).
- [18] A. D. Dolgov, Phys. Rept. **370**, 333 (2002).
- [19] G. B. Gelmini, Phys. Scripta **T121**, 131 (2005).
- [20] R. N. Mohapatra et al., arXiv:hep-ph/0510213.
- [21] R. A. C. Croft, W. Hu and R. Dave, Phys. Rev. Lett. **83**, 1092 (1999).
- [22] M. Tegmark, M. Zaldarriaga and A. J. S. Hamilton, arXiv:hep-ph/0008145.
- [23] X. m. Wang, M. Tegmark and M. Zaldarriaga, Phys. Rev. D **65**, 123001 (2002).
- [24] If so, then there may be Goldstone bosons associated with the broken translational symmetries, which may be very light and hence could mediate new long range forces. Exploring their dynamics could strengthen the bounds on translational symmetry breaking.